



WENSMAN

Genetic Family: W 8195VT2RIB

W 6193RR

94 Day RM

Management Guidelines




Best on stress prone acres



Excels in low yield environments



Monitor late season stalk health south



Agronomic Characteristics

(1 poor : 10 excellent)

<i>Emergence</i>	7
<i>Early Growth</i>	7
<i>GDU's to Silk</i>	1245
<i>Early Flower</i>	6
<i>Brittle Stalks</i>	7

<i>Stalk Strength</i>	7
<i>Root Strength</i>	9
<i>Drought Tolerance</i>	9
<i>GDU's to Black Layer</i>	2320
<i>Black Layers Early</i>	8
<i>Drydown</i>	9
<i>Ear Retention</i>	9
<i>Plant Health</i>	8

<i>Population Tolerance</i>	30000-36000
<i>Corn after Corn</i>	Recommended
<i>Ear Flex</i>	Semi-Flex
<i>Test Weight</i>	8
<i>Kernel Rows</i>	14-18
<i>Cob Color</i>	Red
<i>Shank Length</i>	Medium
<i>Plant Height</i>	Medium
<i>Leaf Angle</i>	Semi-Upright

Disease Tolerance

(1 poor : 10 excellent)

<i>Northern Leaf Blight</i>	7
<i>Southern Leaf Blight</i>	6
<i>Gray Leaf Spot</i>	4
<i>Eyespot</i>	7
<i>Leaf Anthracnose</i>	8
<i>Common Rust</i>	8
<i>Goss's Wilt</i>	6

Soil Recommendations

(1 poor : 10 excellent)

<i>Coarse Soil (sand)</i>	7
<i>Med. Course Soil (silt)</i>	8
<i>Fine Soil (loam)</i>	7
<i>Med. Fine Soil (clay)</i>	6

