

# W3100NR2

## MANAGEMENT GUIDELINES

- Position like W3080NR2 & W3102NR2
- Flexible for use over multiple soils types
- Moves west to east across entire Wensman territory
- IDC tolerance is comparable to W3102NR2


## PRODUCT STRENGTHS

- Goes on every W3102NR2 acre & more
- Rps 1c provides Phytophthora Root Rot protection
- Moderate resistance to Brown Stem Rot
- Travels south well and competes with mid group I material

## PHENOTYPE DESCRIPTION

<b>MATURITY GROUP</b>	1
<b>RELATIVE MATURITY</b>	1.000
<b>VALUE ADDED TRAIT</b>	SCN/GENRR2Y
<b>SOYBEAN CYST NEMATODE</b>	R3, MR14
<b>SOYBEAN CYST NEMATODE GENE</b>	PI 88.788
<b>PHYTOPHTHORA GENE</b>	Rps 1c
<b>FLOWER</b>	Purple
<b>PUBESCENCE</b>	Light Tawny
<b>POD WALL</b>	Brown
<b>HILUM</b>	Black
<b>PLANT TYPE</b>	M/B
<b>PLANT HEIGHT</b>	MT
<b>APPROXIMATE SEED SIZE</b>	Highly Recommended
<b>OIL</b>	M
<b>PROTEIN</b>	M

## NAME OF A MAP



## PRODUCTION

<b>EMERGENCE</b>	8.000
<b>STANDABILITY</b>	8.000
<b>NO-TILL/REDUCED TILL</b>	Excellent
<b>IRON CHLOROSIS (IDC)</b>	7.500
<b>STRESS TOLERANCE</b>	8.000
<b>COARSE SOIL (SANDS)</b>	Excellent
<b>MED COARSE SOIL (SILT)</b>	Excellent
<b>FINE SOIL (LOAM)</b>	Excellent
<b>MED FINE SOIL (CLAY)</b>	RR2

## AGRONOMIC DISEASE

<b>PPR FIELD TOLERANCE</b>	8.000
<b>SCLEROTINA WHITE MOLD (SWM)</b>	7.000
<b>BROWN STEM ROT (BSR)</b>	10.000
<b>SUDDEN DEATH SYNDROME (SDS)</b>	0.000